



MAIKO

LOLIUM PERENNE

Profile

The late, diploid Perennial ryegrass MAIKO shows good yields and high rust resistance. At the same time, MAIKO is one of the few diploid varieties that is suitable for moorland due to its good persistence properties.

National listing/Official recommendation:

DE, NL

Sowing rate: 40 kg/ha

- ✓ diploid variety with moor suitability
- ✓ great rust resistance
- ✓ well balanced yields over all cuts

Suitability

Ploidy	diploid
Maturity	late

Yield characteristics

Dry matter yield total	(1) (2) (3) (4) (5) (6) (7) (8) (9)	medium
Dry matter yield - 1st cut	(1) (2) (3) (4) (5) (6) (7) (8) (9)	medium to high
Dry matter yield - further cuts	(1) (2) (3) (4) (5) (6) (7) (8) (9)	medium

Plant stand development

Early growth development	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input checked="" type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	low to medium
Susceptibility to winter damage	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input checked="" type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	medium
Tendency to lodging	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input checked="" type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	medium
Persistence	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input checked="" type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	medium to high
Sward density	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input checked="" type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	medium to dense

Healthiness

Susceptibility to rust	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input checked="" type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	low to medium
------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------

Quality

Inflorescence formation in aftermath	<input type="radio"/> 1 <input type="radio"/> 2 <input checked="" type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	low
--------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----

Classification according to the descriptive variety list - Bundessortenamt (federal plant variety office, Germany) 2024 and our own results.

All information, recommendations and representations contained herein are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG will not be held liable for any damage or claims for damages resulting from the use of the variety specified in this description. Mixtures may vary if individual varieties are not available. Status 12/2024. Subject to change without prior notice.