

COUNTRY F 2055

Clovergrass, 2-3 years

Two to three years clovergrass mixture for medium to good sites



Profile

This clovergrass mixture has a balanced yield distribution and uniform regrowth behavior. Species were selected to harmonize in competition and persistence and to achieve a good sward density. The high red clover content ensures high protein levels.

- ✓ Balanced yield distribution
- ✓ High yielding
- ✓ High Red clover content



Components

| | |
|-----|--|
| 30% | Festuca pratensis Baltas, Liherold |
| 30% | Trifolium pratense Milvus, Taifun |
| 20% | Lolium perenne (intermediate) Explosion |
| 20% | Phleum pratense Lischka |

If individual varieties are not available, they are replaced by equivalent ones.

Locations

| | |
|-----------|---|
| Dry | <div><div>+</div><div>+</div><div></div></div> |
| Normal | <div><div>+</div><div>+</div><div>+</div></div> |
| Wet | <div><div>+</div><div>+</div><div>+</div></div> |
| Peat Soil | <div><div>+</div><div>+</div><div></div></div> |
| Altitude | <div><div>+</div><div>+</div><div></div></div> |

+++ = highly suitable | ++ = suitable | + = conditionally suitable | - = not given

Cultivation notes

| | |
|----------------|---------------|
| Sowing rate | 20 - 25 kg/ha |
| New sowing | 20 - 25 kg/ha |
| Usage per year | 3 - 4 |

All specified informations, recommendations and representations mentioned here are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG excludes adhesion for damage or claims for damages, resulting of the use for the variety specified in this description. Mixture compositions may change if individual varieties are not available. As of 07/2023. Subject to change without notice.