



COUNTRY G 2004

With clover

Top performing mixture with clover for intensive grassland

Profile

Widely usable mixture with uniform yield distribution by using all maturity groups. White clover increases the protein yield and provides N-fixation.

- ✓ Stable yield distribution
- ✓ English ryegrass of all maturity groups
- ✓ White clover for N-fixation



Components

| | |
|-----|---|
| 45% | Lolium perenne (late) Chevalier, Valerio |
| 25% | Lolium perenne (intermediate) AstonHockey, Botond |
| 25% | Lolium perenne (early) Mirtello |
| 5% | Trifolium repens Bianca, Liflex |

If individual varieties are not available, they are replaced by equivalent ones.

Sowing pattern

New sowing, Overseeding

Locations

| | |
|-----------|-------|
| Dry | ⊕ ○ ○ |
| Normal | ⊕ ⊕ ○ |
| Wet | ⊕ ⊕ ⊕ |
| Peat Soil | ⊕ ○ ○ |
| Altitude | ⊕ ⊕ ○ |

Usage

| | |
|---------------------|-------|
| Grazing | ⊕ ⊕ ⊕ |
| Grazing and cutting | ⊕ ⊕ ⊕ |
| Cutting | ⊕ ⊕ ⊕ |

+++ = highly suitable | ++ = suitable | + = conditionally suitable | - = not given

Cultivation notes

| | |
|-------------------------------------|---------------|
| New sowing | 40 kg/ha |
| Overseeding | 20 - 25 kg/ha |
| Overseeding, several times per year | 7 - 10 kg/ha |
| Usage per year | 4 - 5 |

All information, recommendations and representations contained herein are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG will not be held liable for any damage or claims for damages resulting from the use of the variety specified in this description. Mixtures may vary if individual varieties are not available. Status 12/2024. Subject to change without prior notice.