COUNTRY GRASSLAND 2441 ORGANIC



New sowing without clover

Profile

The different grass species provide a dense sward and palatable forage, especially for horses. The special selection of perennial ryegrass, tall fescue and red fescue withstand the horses deep browsing and hard treading. This mixture is suitable for grazing and cutting, as well as for outdoor chicken runs.

- ✓ Robust yielding mixture
- ✓ Winterhardy
- ✓ High sward density

Components

| 30% | Lolium perenne (intermediate) Explosion |
|-----|---|
| 20% | Festuca arundinacea Lipalma |
| 15% | Festuca pratensis Liherold |
| 13% | Festulolium Fedoro |
| 12% | Phleum pratense Aturo |
| 10% | Festuca rubra Rafael |



Sowing pattern

New sowing

Locations

| Dry | | | |
|------------------------------|--|--|--|
| Normal | + + | | |
| Wet | + + | | |
| Peat Soil | • • | | |
| Altitude | | | |
| Usage | | | |
| Grazing | | | |
| Grazing and cutting | + + + | | |
| Cutting | + + | | |
| Extensive usage | + | | |
| Fodder values | | | |
| Energy density | | | |
| Digestibility | + + | | |
| Protein | + | | |
| +++ = highly suitable ++ = | +++ = highly suitable ++ = suitable + = conditionally suitable - = not given | | |



Cultivation notes

| New sowing | 40 kg/ha |
|----------------|-------------------------|
| Sowing time | from March to September |
| Usage per year | 2 |

All organic mixtures are produced with 100 % organic content, if possible, in accordance with the EU Organic Farming Regulation 2018/848. For mixtures with at least 70% organic content, the farmer must obtain approval from his inspection body for the use of the conventional content before sowing, starting with production in January 2022. We aim to use at least 2 varieties per ripening group for mix proportions of more than 20 % of one species and for German ryegrass. The choice of varieties may vary depending on production. If individual species or varieties are not available, they are replaced by equivalent ones. The seed strengths are calculated for an optimal seedbed; in case of poor external conditions - such as insufficient seedbed, late sowing date, insufficient nutrient supply and unfavourable weather - the seed strengths should be adjusted upwards.

All specified informations, recommendations and representations mentioned here are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG excludes adhesion for damage or claims for damages, resulting of the use for the variety specified in this description. Mixture compositions may change if individual varieties are not available. As of 07/2023. Subject to change without notice.