



COUNTRY FIELD FORAGE 2488 ORGANIC

Clover grass intercropping

Profile

COUNTRY Field forage 2488 is a composition of compatible annual ryegrass and clovers. This mixture is fast growing, high yielding and palatable. It meets the demands concerning green fodder as well as silage.

- ✓ Fast growing through Annual ryegrasses
- ✓ For intensive use
- ✓ High yielding and high quality forage



Components

70%	Lolium westerwoldicum Alberto*, Arnoldo*
15%	Egyptian clover
15%	Persian clover

* also suitable for intercropping, yield in 1st cut above average

Sowing pattern

New sowing

Locations

Dry	<div><div>+</div><div></div><div></div></div>
Normal	<div><div>+</div><div>+</div><div>+</div></div>
Wet	<div><div>+</div><div>+</div><div>+</div></div>
Peat Soil	<div><div>+</div><div></div><div></div></div>
Altitude	<div><div>+</div><div>+</div><div></div></div>

Usage

Grazing	<div><div>+</div><div>+</div><div></div></div>
Grazing and cutting	<div><div>+</div><div></div><div></div></div>
Cutting	<div><div>+</div><div>+</div><div>+</div></div>
Extensive usage	<div><div>+</div><div>+</div><div></div></div>

Fodder values

Energy density	<div><div>+</div><div>+</div><div>+</div></div>
Digestibility	<div><div>+</div><div>+</div><div>+</div></div>
Protein	<div><div>+</div><div>+</div><div>+</div></div>

+++ = highly suitable | ++ = suitable | + = conditionally suitable | - = not given

Cultivation notes

New sowing	35 - 40 kg/ha
Sowing time	from July to August
Usage per year	1 - 2

All organic mixtures are produced with 100 % organic content, if possible, in accordance with the EU Organic Farming Regulation 2018/848. For mixtures with at least 70% organic content, the farmer must obtain approval from his inspection body for the use of the conventional content before sowing, starting with production in January 2022. We aim to use at least 2 varieties per ripening group for mix proportions of more than 20 % of one species and for German ryegrass. The choice of varieties may vary depending on production. If individual species or varieties are not available, they are replaced by equivalent ones. The seed strengths are calculated for an optimal seedbed; in case of poor external conditions - such as insufficient seedbed, late sowing date, insufficient nutrient supply and unfavourable weather - the seed strengths should be adjusted upwards.

All information, recommendations and representations contained herein are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG will not be held liable for any damage or claims for damages resulting from the use of the variety specified in this description. Mixtures may vary if individual varieties are not available. Status 12/2024. Subject to change without prior notice.