



FRIEDA

WHITE LUPIN

A real protein bomb

Profile

FRIEDA is a real protein miracle. In contrast to soya, it has a significantly better climate adaptation, a lower heat requirement and can thus be grown on many sites. Due to its unique anthracnose tolerance, it provides farmers with high yield security. FRIEDA is suitable for animal feeding.

Distribution in: AT, CH, DE, NL, UK • Breeder: LLA Triesdorf

- ✓ Branching type with low content of bitter substances
- ✓ High protein yield for feed use
- ✓ Excellent drought tolerance due to taproot
- ✓ Tolerance against anthracnose
- ✓ Very good threshing ability due to homogenous grain/straw ripening and high core build-up
- ✓ Better yield potential, higher protein content with better space stability than blue lupins



Agronomic properties

Determinate growth	<input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	missing
Beginning of flowering	<input type="radio"/> 1 <input type="radio"/> 2 <input checked="" type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	early
Ripening	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input checked="" type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	early to inter
Plant height	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input checked="" type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	medium
Bitterness level	<input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	low in bitter substances

Tends to

Lodging	<input type="radio"/> 1 <input type="radio"/> 2 <input checked="" type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	low
---------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----

Yield characteristics

Thousand grain mass	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input checked="" type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	high
Grain yield	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input checked="" type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	medium to high
Gross protein yield	<input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input checked="" type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	high

Classification according to the descriptive variety list - Bundessortenamt (federal plant variety office, Germany) 2025 and our own results.

Quality

Crude protein content	<input type="radio"/> 1 <input type="radio"/> 2 <input checked="" type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9	low
-----------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----

Locations

All soils except waterlogging

Special characteristics

Greater yield security through tolerance to anthracnose, the most important lupin disease, which can lead to a complete loss of yield.

Sowing

60 grains/m² for row distances of 12-25 cm

Sowing date

Mid of March - mid of April

Usage

High-quality protein feed for cattle feeding

All information, recommendations and representations contained herein are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG will not be held liable for any damage or claims for damages resulting from the use of the variety specified in this description. Mixtures may vary if individual varieties are not available. Status 12/2024. Subject to change without prior notice.