

Profile

The species richness of BetaSola ensures good soil structure and thereby differs considerably from the cultivation of a single nematode-resistant monocrop. The combination of species has a wide spectrum of activity. In this way, the various nematode-resistant oil radish varieties help to reduce beet nematodes (Heterodera schachtii) and Trichodorus. Furthermore, multi-resistant oil radish is resistant to rootknot nematodes. The mix partner bristle oat also reduces soil nematodes (Pratylenchus).

- ✓ Nematode reduction & soil protection
- ✓ Wide action spectrum (Heterodera schachtii and Trichodorus)
- ✓ Specially adapted to potato and beet production technology

N-Potential approx. 60 kg/ha

Components

21.1%	Legumes
33.7%	Cruciferous
	Oil radish, Common vetch, Bristle oat, Niger, Egyptian clover
Seed shares (%)	

Sowing pattern

blank seed, mulch sowing, direct sowing, spreader device on tillage implement, drone

Crop rotations

Potatoes, Sugar beet, Cereals, Maize

Cultivation notes

Sowing rate	35 - 40 kg/ha
Sowing time	from middle of July to end of August
Usage	Cover crop, summer intercrop, winter intercrop, green fertilization, Extensive grazing, normal sowing time, N-Potential: approx. 60 kg/ha, medium to low proportion of legumes



All information, recommendations and representations contained herein are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG will not be held liable for any damage or claims for damages resulting from the use of the variety specified in this description. Mixtures may vary if individual varieties are not available. Status 12/2024. Subject to change without prior notice.

