



FRIEDA ORCANIC

White lupin organic

A real protein bomb

Profile

FRIEDA is a real protein miracle. In contrast to soya, it has a significantly better climate adaptation, a lower heat requirement and can thus be grown on many sites. Due to its unique anthracnose tolerance, it provides farmers with high yield security. FRIEDA is suitable for animal feed; its use in human nutrition is still being tested.

Distribution in: AT, CH, DE • Breeder: LLA Triesdorf

- ✓ Branching type with low content of bitter substances
- ✓ High protein yield for feed use
- ✓ Also usable for human nutrition due to lower alkaloid content
- ✓ Tolerance against anthracnose
- ✓ Better yield potential, higher protein content with better space stability than blue lupins
- ✓ Deep-rooted taproot and therefore very insensitive to summer dryness
- ✓ Very good threshing ability due to homogenous grain/straw ripening and high core build-up

Agronomic properties

| | | |
|------------------------|--|--------------------------|
| Determinate growth | <input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | missing |
| Beginning of flowering | <input type="radio"/> 1 <input type="radio"/> 2 <input checked="" type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | early |
| Ripening | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input checked="" type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | early to inter |
| Plant height | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input checked="" type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | medium |
| Bitterness level | <input checked="" type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | low in bitter substances |

Tends to

| | | |
|---------|--|-----|
| Lodging | <input type="radio"/> 1 <input type="radio"/> 2 <input checked="" type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | low |
|---------|--|-----|

Yield characteristics

| | | |
|---------------------|--|----------------|
| Thousand grain mass | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input checked="" type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | high |
| Grain yield | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input checked="" type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | medium to high |
| Gross protein yield | <input type="radio"/> 1 <input type="radio"/> 2 <input type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input checked="" type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | high |

Classification according to the descriptive variety list - Bundessortenamt (federal plant variety office, Germany) 2024 and our own results.

Quality

| | | |
|-----------------------|--|-----|
| Crude protein content | <input type="radio"/> 1 <input type="radio"/> 2 <input checked="" type="radio"/> 3 <input type="radio"/> 4 <input type="radio"/> 5 <input type="radio"/> 6 <input type="radio"/> 7 <input type="radio"/> 8 <input type="radio"/> 9 | low |
|-----------------------|--|-----|

Locations

All soils except waterlogging

Special characteristics

Greater yield security through tolerance to anthracnosis, the most important lupin disease, which can lead to a complete loss of yield.

Sowing

60 grains/m² for row distances of 12-25 cm

Sowing date

Mid of March - mid of April

Usage

High-quality protein feed for cattle feeding in organic quality

All information, recommendations and representations contained herein are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG will not be held liable for any damage or claims for damages resulting from the use of the variety specified in this description. Mixtures may vary if individual varieties are not available. Status 12/2024. Subject to change without prior notice.