

### Profile

PONANT is a sorghum hybrid with a high yield potential. Approximately 85 to 95 days after sowing, the early hybrid starts to flowering. The half-open risp has orange to light-brown grains with a low tannin content. Furthermore, the variety has a good health status and is stable. The grain sorghum PONANT is used as fodder as well as for biogas or animal nutrition.

- ✓ Early-ripening grain sorghum
- ✓ Orange-brownish grains
- ✓ Low tannin content

# Usage

Silo	
------	--

Dualtype

## Sowing

Sowing rate	25 - 35 Plants/m²
Seed distance	30 - 70 cm

## Sowing date

At a soil temperature of 12 - 14°C (mid-May - early June).

#### Seedbed

An optimal seed bed for quick warming and emergence of the sorghum plants.

## Seeding procedure

Sorghum is sown in rows with an beet or corn sowing machine with sorghum discs. As an alternative sorghum can also be broadcast-sown. When selecting your sowing machine, please take in mind that seed sizes might vary greatly amongst varieties.



# Sowing depth

2-4 cm

### **Fertilization**

Nitrogen: 60 - 80 kg/haPhosphate: 35 - 45 kg/haPotassium: 120 - 150 kg/ha

## Plant protection

A preparation of a false seed bed before sowing reduces weed pressure during initial development. A combination of mechanical and chemical weed control, whenever possible, is also recommended.

All information, recommendations and representations contained herein are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG will not be held liable for any damage or claims for damages resulting from the use of the variety specified in this description. Mixtures may vary if individual varieties are not available. Status 12/2024. Subject to change without prior notice.

